



VETERINARY GENETICS LABORATORY
SCHOOL OF VETERINARY MEDICINE
ONE SHIELDS AVENUE
DAVIS, CALIFORNIA 95616-8744

TELEPHONE: (530) 752-2211
FAX: (530) 752-3556

BENGAL PRA TEST REPORT

REBECCA BÖSCH HAUPTSTRASSE 84 9434 AU SWITZERLAND	Case: CAT107017 Date Received: 28-Jun-2018 Print Date: 29-Jun-2018 Report ID: 6489-8609-2012-5146 Verify report at www.vgl.ucdavis.edu/myvgl/verify.html
Cat: OCTOPUSSY DES GRIFFES DE FEU <i>Reg:</i> DOB: 02/24/2018 Sex: Female Breed: Bengal Microchip: 250268731916961 Color: Seal mink spotted tabby	
Sire: KAMISHA HOODOO SKY <i>Reg:</i> 977200008809132 Dam: MASCARA DES GRIFFES DE FEU <i>Reg:</i> 250268731630299	

BENGAL PRA RESULT

N/N

Result Codes:

- N/N Normal - no copies of the PRA-b mutation.
- N/PRA Carrier - 1 copy of the PRA-b mutation; vision will be normal.
- PRA/PRA Affected - 2 copies of the PRA-b mutation; cat will develop clinical signs of Bengal PRA.

For more detailed information on Bengal PRA results, please go to:
www.vgl.ucdavis.edu/services/cat/BengalPRA.php

BENGAL PROGRESIVE RETINAL ATROPHY

The Veterinary Genetics Laboratory at the University of California – Davis and the Lyons Feline Genetics and Comparative Medicine Laboratory at the University of Missouri announce the release of the **new genetic test for progressive retinal atrophy (PRA)**, which causes an autosomal recessive blindness in Bengal cats. The disease causes the destruction of the cells that register light (photoreceptors) in the back of the eye (the retina). The loss of the cells begins around 7 weeks of age and slowly progresses until the cat has very compromised vision by approximately 2 years of age¹. However, blindness develops at different rates in different cats. We have examples of cats over 2 years of age chasing a laser pointer; however vision testing by an ophthalmologist indicated that the cats should be blind. Blind cats tend to have more difficulty at night, sometimes becoming more vocal and more attached to their owners. The pupils are usually more dilated for affected cats than for cats with normal vision in the same lighting conditions. Affected cats also tend to carry their whiskers in a more forward position. Once affected cats know their surroundings, they are very mobile and active. Our thanks to the breeders who came forward and helped us establish a colony so that we could define the condition and find the gene responsible for this defect.

The mutant DNA variant appears to be novel to the Bengal breed and occurred early in a popular lineage of the Bengals. We expect Bengal cats worldwide to have the condition and we have had reports of affected cats in the United Kingdom, Europe and the USA. Bengal PRA is autosomal recessive, thus two copies of the mutant DNA variant are required for the cats to be blind. The blindness can be detected either by the DNA test or by an eye exam prior breeding age. Carriers, cats with one copy of the mutation, can only be detected by the DNA test.

Because the mutation is recent in evolutionary time, an extended DNA region and several genes around the casual mutation had to be examined as many DNA variants appeared to be perfectly linked with the blindness. This is a common and well known phenomenon, called linkage disequilibrium. Other examples of linkage disequilibrium have been published for cat DNA variants that cause *Brown* coloration² and cat AB blood group.³

Several DNA variants within the region needed to be evaluated and after working with European breeders, many non-causal variants were eliminated. Initially, a gene mutation in a known vision gene was implicated, but after screening hundreds of cats, a second mutation in a different and novel vision gene was identified as being perfectly

